

## Intravenous (IV) Fluids

IV fluids are utilized for resuscitation, to replace fluid losses, and for maintenance hydration in patients with contraindications to oral fluid intake. Like all prescribed medications, IV fluids have benefits and risks. Fluid therapy should be carefully administered, with specific goals or endpoints established to avoid inadvertent fluid overload or acid-base alterations.

### Crystalloids

Crystalloid solutions contain small molecules that flow easily across semipermeable membranes, from the bloodstream into the cells and body tissues. Crystalloid solutions are distinguished by the relative tonicity (before infusion) in relation to plasma and are categorized as isotonic, hypotonic, or hypertonic.

#### **Isotonic solutions**

Isotonic solutions have a concentration of dissolved particles similar to plasma and an osmolality of 250 to 375 mOsm/L. These fluids remain within the extracellular compartment and are distributed between intravascular (blood vessels) and interstitial (tissue) spaces, increasing intravascular volume. They are used primarily to treat fluid volume deficit.

#### *General nursing considerations:*

- Document baseline vital signs, body weight, lung sounds, heart sounds, and note presence or absence of edema. Continue monitoring during and after the infusion.
- Monitor for continued signs of hypovolemia, including urine output less than 0.5 mL/kg/hour, poor skin turgor, tachycardia, weak pulse, and hypotension.
- Monitor for signs of hypervolemia such as hypertension, bounding pulse, pulmonary crackles, dyspnea, shortness of breath, peripheral edema, weight gain, jugular vein distension (JVD) and extra heart sounds such as S3.
- A daily serum metabolic panel is recommended for patients receiving continuous IV fluids to monitor sodium, potassium, and glucose levels, as well as renal function.
- More frequent metabolic panel analysis is indicated for unstable patients or those with significant electrolyte or acid-base disturbances.

I.V. Fluid Osmolarity Composition	Uses/Clinical Considerations
<p><b>0.9% NaCl</b> (Normal Saline Solution, NSS)</p> <p>308 mOsm/L Na<sup>+</sup> 154 mmol/L Cl<sup>-</sup> 154 mmol/L</p>	<ul style="list-style-type: none"> <li>• <b>Often the default choice for resuscitation due to availability, low cost, and compatibility with blood products and most medications.</b></li> <li>• Used to replace fluid loss from hemorrhage, severe vomiting or diarrhea, heavy drainage from GI suction, fistulas or wounds.</li> <li>• Use to treat shock, mild hyponatremia, and hypercalcemia.</li> <li>• Caution in cardiac or renal disease.</li> <li>• Monitor for hyperchloremic acidosis with large volumes of fluid replacement with 0.9% NaCl.</li> </ul>
<p><b>Lactated Ringer's Solution</b> (LR, Ringer's Lactate)</p> <p>273 mOsm/L Na<sup>+</sup> 130mEq/L K<sup>+</sup> 4 mEq/L Ca<sup>++</sup> 3 mEq/L Cl<sup>-</sup> 109 mEq/L</p>	<ul style="list-style-type: none"> <li>• More closely matches plasma chemistry. For this reason, LR and plasmalyte are known as "balanced" solutions.</li> <li>• <b>Balanced crystalloids are preferred for major volume resuscitation</b> to mitigate the risk of acidosis and acute kidney injury.</li> <li>• Used to treat shock, acute blood loss, hypovolemia due to burns, third-space fluid shift, GI loss and fistula drainage, electrolyte loss, and metabolic acidosis.</li> <li>• Use with caution in patients who cannot metabolize lactate, (e.g., liver disease).</li> <li>• Do not administer if pH is greater than 7.5. (Normal liver will convert LR to bicarbonate, worsening alkalosis).</li> <li>• Avoid in patients with hyperkalemia or severe renal failure (LR contains potassium).</li> </ul>
<p><b>5% dextrose in water</b> (D<sub>5</sub>W)</p> <p>253 mOsm/L 5 g dextrose/100mL 50 g dextrose/L 170 calories/L</p>	<ul style="list-style-type: none"> <li>• Both isotonic and hypotonic. Initially dilutes osmolality of extracellular fluid (hypotonic); once cell has used dextrose, remaining saline and electrolytes act isotonic, expanding the extracellular compartment.</li> <li>• Provides free water for the kidneys, aiding renal excretion of solutes.</li> <li>• May be used to treat hypernatremia.</li> <li>• Should not be used alone to treat fluid volume deficit because it dilutes plasma electrolyte concentrations.</li> <li>• Contraindicated in resuscitation, early postoperative period, and patients with known or suspected increased intracranial pressure (ICP).</li> <li>• Provides some calories, but not enough nutrition for prolonged use.</li> </ul>
<p><b>Additional isotonic solutions:</b></p>	
<p><b>Ringer's Solution</b> Similar to LR but does not contain lactate. Not an alkalizing agent; not ideal for patients with metabolic acidosis.</p>	

**PlasmaLyte**

Electrolyte composition similar to plasma; can be infused with packed red blood cells. Less likely than other fluids to lead to dilutional or hyperchloremic acidosis.

**Hypotonic solutions**

Hypotonic solutions have a concentration of dissolved particles lower compared to plasma and an osmolality less than 250 mOsm/L. Hypotonic fluids lower serum osmolality within the vascular space by causing fluid to shift out of the blood into the cells and tissue spaces.

Typically used to treat conditions causing intracellular dehydration, such as diabetic ketoacidosis and hyperosmolar hyperglycemic states.

*General nursing considerations:*

- May worsen existing hypovolemia and hypotension causing cardiovascular collapse.
- Monitor for signs of fluid volume deficit, such as confusion in older adults and dizziness. Never administer to patients at risk for increased ICP as the potential fluid shift may cause cerebral edema. Avoid in patients with liver disease, trauma or burns.

I.V. Fluid Osmolarity Composition	Uses/Clinical Considerations
<p><b>0.45% NaCl</b> (1/2 NS)</p> <p>154 mOsm/L Na<sup>+</sup> 77 mEq/L Cl<sup>-</sup> 77 mEq/L</p>	<ul style="list-style-type: none"> <li>• Used to treat hyponatremia.</li> <li>• Caution in patients with heart failure, severe renal insufficiency, and edema with sodium retention.</li> <li>• May cause fluid overload resulting in decreased electrolyte concentrations, congested states or pulmonary edema.</li> <li>• Rapid infusion may cause hemolysis of red blood cells (RBCs).</li> </ul>
<b>Additional hypotonic solutions:</b>	
<p><b>0.33% NaCl</b> Allows kidneys to retain needed amounts of water. Caution in patient with heart failure and severe renal insufficiency. Adverse effects include pulmonary edema, febrile reactions. Typically administered with dextrose to increase tonicity.</p>	
<p><b>0.225% NaCl</b> The most hypotonic fluid available, often recommended as maintenance fluid for pediatric patients. Avoid rapid infusion to prevent hemolysis. Avoid use unless mixed with dextrose.</p>	
<p><b>2.5% dextrose in water (D<sub>2.5</sub>W)</b> Used to treat dehydration and decrease sodium and potassium levels. Not administered with blood as it can cause hemolysis of RBCs.</p>	

**Hypertonic solutions**

Hypertonic solutions have a concentration of dissolved particles higher than plasma and an osmolality greater than 375 mOsm/L. A higher solute concentration causes the osmotic pressure gradient to draw water out of cells, increasing extracellular volume. These fluids are often used as volume expanders and may be prescribed for hyponatremia (low sodium). They may also benefit patients with cerebral edema.

*General nursing considerations:*

- Administer only in high-acuity areas.
- For short-term use to correct critical electrolyte abnormalities.
- Monitor electrolytes and assess for hypervolemia. May cause fluid volume overload and pulmonary edema.
- Avoid in patients with cardiac or renal conditions who are dehydrated, and in patients with diabetic ketoacidosis.

I.V. Fluid Osmolarity Composition	Uses/Clinical Considerations
<p><b>3% NaCl</b> 1030 mOsm/L ----- <b>5% NaCl</b> 1710 mOsm/L</p>	<ul style="list-style-type: none"> <li>• Used for treatment of severe, critical symptomatic hyponatremia and cerebral edema.</li> <li>• Give slowly and cautiously to avoid intravascular fluid volume overload, pulmonary edema, and the rare life-threatening complication of central pontine myelinolysis.</li> <li>• Rapid infusion may cause hypernatremia or hyperchloremia.</li> </ul>
<p><b>Dextrose 5% in 0.45% NaCl</b> (D<sub>5</sub> ½ NS)  406 mOsm/L</p>	<ul style="list-style-type: none"> <li>• Used as maintenance IV fluid for patients unable to ingest adequate fluids orally.</li> <li>• Monitor closely for fluid volume overload.</li> <li>• Monitor blood glucose, sodium, and potassium levels.</li> </ul>
<p><b>Dextrose 5% in 0.9% NaCl</b> (D<sub>5</sub>NS)  560 mOsm/L</p>	<ul style="list-style-type: none"> <li>• Provides calories, water and electrolytes</li> <li>• Monitor closely for fluid volume overload and pulmonary edema.</li> </ul>
<p><b>Dextrose 5% in Lactated Ringer's</b> (D<sub>5</sub>LR)  527 mOsm/L</p>	<ul style="list-style-type: none"> <li>• Provides calories, water and electrolytes</li> <li>• Contains sodium lactate which may be used to treat metabolic acidosis.</li> </ul>
<p><b>10% Dextrose in water</b> (D<sub>10</sub>W)  505 mOsm/L 10 g dextrose/100mL 340 calories/L</p>	<ul style="list-style-type: none"> <li>• Provides free water and calories, but no electrolytes.</li> <li>• Contraindicated in patients with intracranial or intra-spinal hemorrhage, delirium tremens, severe dehydration, anuria, hepatic coma.</li> <li>• Use central line if possible. Do not infuse through same IV line as blood products due to possibility of RBC hemolysis.</li> <li>• Monitor blood glucose closely. Use with caution in patients with diabetes mellitus. Monitor for hypokalemia.</li> <li>• May cause phlebitis, vein damage and thrombosis at the injection site.</li> </ul>

	<ul style="list-style-type: none"> <li>• Rapid infusion may cause diuresis, hyperglycemia, glycosuria,</li> <li>• hyperosmolar syndrome (mental confusion, loss of consciousness), fluid and/or solute overload, overhydration, or pulmonary edema.</li> </ul>
<b>Additional hypertonic solutions:</b>	
<b>20% Dextrose in water (D<sub>20</sub>W):</b> Acts as an osmotic diuretic, causes a fluid shift between various compartments. Promotes diuresis.	
<b>50% Dextrose in water (D<sub>50</sub>W):</b> Administered via IV bolus to treat patients with severe hypoglycemia.	

### Colloids

- Colloid solutions contain large molecules that do not pass through semipermeable membranes and therefore remain in the blood vessels. Also known as volume/plasma expanders, colloids expand intravascular volume by drawing fluid from the interstitial space into the vessels through higher oncotic pressure. Colloids (usually albumin) are not used as first-line agents for general fluid resuscitation. Colloids are significantly more expensive than crystalloids, and synthetic colloids carry major risks. Therefore, unless a specific indication exists, safety and cost-effectiveness favor crystalloids.

#### General nursing considerations:

- Before administering a colloid, take a careful allergy history.
- Use 18-gauge or larger needle for administration of colloid solutions.
- Monitor intake and output closely and for signs of hypervolemia: hypertension, dyspnea, crackles in lungs, jugular venous distension, edema, bounding pulse, and extra heart sounds such as S3.
- Monitor coagulation indexes.

I.V. Fluid Osmolarity	Clinical Considerations
<b>Albumin (5%)</b>  309 mOsm/L ----- <b>Albumin (25%)</b>  312 mOsm/L	<ul style="list-style-type: none"> <li>• Human albumin solution</li> <li>• Used for hypotension in patients with spontaneous bacterial peritonitis and during large volume paracentesis in patients with cirrhosis.</li> <li>• For adults with septic shock who require substantial amounts of crystalloid to maintain mean arterial pressure (MAP) over 65, the Surviving Sepsis Campaign Guidelines suggest adding albumin to the fluid resuscitation regimen.</li> <li>• Monitor for an allergic reaction or anaphylaxis.</li> <li>• Contraindicated in severe anemia, heart failure or known sensitivity to albumin.</li> </ul>

**Additional Colloid solutions (rarely used):**

**Low-molecular weight dextran (LMWD); High-molecular weight dextran (HMWD)**

Contains polysaccharide molecules that behave like colloids. Dextran carries several significant risks that make them less favorable than newer solutions, including coagulopathy, Acute Kidney Injury (AKI), and anaphylaxis.

Dextran is sometimes still used post-operatively, particularly in vascular or reconstructive surgery, for its properties that inhibit red blood cell clumping and reduce blood viscosity.

**Hydroxyethyl Starch (HES), Hetastarch**

Once common but now heavily restricted or contraindicated due to proven risks. Strongly contraindicated in patients with sepsis, severe trauma, or burn injury due to evidence linking its use to increased risk of Acute Kidney Injury and death. Many international guidelines have removed these colloids from routine use.

**References:**

Barlow, A., Barlow, B., Tang, N., Shah, B., King, A. (2020). Intravenous Fluid Management in Critically Ill Adults: A Review. *Crit Care Nurse* 1. 40 (6): e17–e27. <https://doi.org/10.4037/ccn2020337>

Evans, L., Rhodes, A., Alhazzani, W., Antonelli, M., Coopersmith, C. M., French, C., Machado, F. R., McIntyre, L., Ostermann, M., Prescott, H. C., Schorr, C., Simpson, S., Wiersinga, W. J., Alshamsi, F., Angus, D. C., Arabi, Y., Azevedo, L. C. P., Backer, D. D., Behrendt, J., et al. (2021). Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock 2021. *Intensive Care Medicine*, 47(11), 1181–1247. <https://doi.org/10.1007/s00134-021-06506-y>

Guest M. (2020). Understanding the principles and aims of intravenous fluid therapy. *Nursing Standard*. <https://doi.org/10.7748/ns.2020.e11459>

Gross, W., Samarin, M., & Kimmons, L. (2017). Choice of Fluids for Resuscitation of the Critically Ill: What Nurses Need to Know. *Critical Care Nursing Quarterly*, 40(4), 309-322. <https://doi.org/10.1097/CNQ.0000000000000170>

Malbrain, M.L.N.G., Langer, T., Annane, D., Gattinoni, L., Elbers, P., Hahn, R.G., De Laet, I., Minini, A., Wong, A., Ince, C., Muckart, D., Mythen, M., Caironi, P., Van Regenmortel, N. (2020). Intravenous fluid therapy in the perioperative and critical care setting: Executive summary of the International Fluid Academy (IFA). *Annals of Intensive Care*. 10(1),64. <https://doi.org/10.1186/s13613-020-00679-3>.

Semler, M. W., Self, W. H., Wanderer, J. P., Ehrenfeld, J. M., Wang, L., Byrne, D. W., Stollings, J. L., Kumar, A. B., Hughes, C. G., Hernandez, A., Guillaumondegui, O. D., May, A. K., Weavind, L., Casey, J. D., Siew, E. D., Shaw, A. D., Bernard, G. R., Rice, T. W., & SMART Investigators and the Pragmatic Critical Care Research Group (2018). Balanced Crystalloids versus Saline in Critically Ill Adults. *The New England journal of medicine*, 378(9), 829–839. <https://doi.org/10.1056/NEJMoa1711584>

Sterns, R. (2025, July 18). Maintenance and replacement fluid therapy in adults. *UpToDate*. <https://www.uptodate.com/contents/maintenance-and-replacement-fluid-therapy-in-adults>